

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



Application Number: 10/631,029  
Filing Date: 07/29/2003  
First Named Inventor: Rajinder Singh  
Attorney Docket Number: 064311-1602  
Identifier: F6

(43) International Publication Date  
5 April 2001 (05.04.2001)

PCT

(10) International Publication Number  
**WO 01/23362 A2**

(51) International Patent Classification<sup>7</sup>: **C07D 239/48**,  
251/18, 239/50, 403/12, 521/00, 405/14, A61K 31/505,  
A61P 35/00

Dannstadt (DE). **ROSENBERG, Jörg** [DE/DE]; Bruch-  
strasse 29, 67158 Ellerstadt (DE). **BREITENBACH, Jörg**  
[DE/DE]; Hans-Sachs-Ring 95A, 68199 Mannheim (DE).

(21) International Application Number: PCT/EP00/09149

(74) Agent: **GOLDSCHIED, Bettina**; BASF Aktiengesellschaft, 67056 Ludwigshafen (DE).

(22) International Filing Date:  
19 September 2000 (19.09.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
199 45 982.7 24 September 1999 (24.09.1999) DE

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

(71) Applicant (*for all designated States except US*): **KNOLL AKTIENGESELLSCHAFT** [DE/DE]; 67061 Ludwigshafen (DE).

**Published:**


— Without international search report and to be republished upon receipt of that report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **HANTKE, Thomas** [DE/DE]; Landauerstrasse 5, 67067 Ludwigshafen (DE). **REHBOCK, Bettina** [DE/DE]; Wasgauring 28, 67125

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: RATE-CONTROLLED PARTICLES



(57) Abstract: Rate-controlled particles, comprising compounds of formula (I) as a solid dispersion.

WO 01/23362 A2

Rate-controlled particles


Specification

5

The present invention concerns pharmaceutical compositions in the form of rate-controlled particles, comprising compounds of the formula (I) to (VI)

10 (I) is an antiviral compound of formula

15



a N-oxide, a pharmaceutically acceptable addition salt or a  
20 stereochemically isomeric form thereof, wherein

Y is CR<sup>5</sup> or N;

A is CH, CR<sup>4</sup> or N;

n is 0, 1, 2, 3 or 4;

25 Q is -NR<sup>1</sup>R<sup>2</sup> or when Y is CR<sup>5</sup> then Q may also be hydrogen;

R<sup>1</sup> and R<sup>2</sup> are each independently selected from hydrogen, hydroxy, C<sub>1-12</sub>alkyl, C<sub>1-12</sub>alkyloxy, C<sub>1-12</sub>alkylcarbonyl, C<sub>1-12</sub>alkyloxycarbonyl, aryl, amino, mono- or di(C<sub>1-12</sub>alkyl)-amino, mono- or di(C<sub>1-12</sub>alkyl)aminocarbonyl wherein each of  
30 the aforementioned C<sub>1-12</sub>alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxy-C<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, aminocarbonyl, aminocarbonylamino, mono- or  
35 di(C<sub>1-6</sub>alkyl)amino, aryl and Het; or

R<sup>1</sup> and R<sup>2</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-12</sub>alkyl)aminoC<sub>1-4</sub>-alkylidene;

R<sup>3</sup> is hydrogen, aryl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy-carbonyl, C<sub>1-6</sub>alkyl substituted with C<sub>1-6</sub>alkyloxycarbonyl; and  
40 each R<sup>4</sup> independently is hydroxy, halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalo-methyloxy, or when Y is CR<sup>5</sup> then R<sup>4</sup> may also represent C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl;

45 R<sup>5</sup> is hydrogen or C<sub>1-4</sub>alkyl;

L is -X<sup>1</sup>-R<sup>6</sup> or -X<sup>2</sup>-Alk-R<sup>7</sup> wherein


- 5**  $R^6$  and  $R^7$  each independently are phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkyloxycarbonyl, formyl, cyano, nitro, amino, and trifluoromethyl; or when  $Y$  is  $CR^5$  then  $R^6$  and  $R^7$  may also be selected from phenyl substituted with one, two, three, four or five substituents each independently selected from aminocarbonyl, trihalomethyloxy and trihalomethyl; or when  $Y$  is  $N$  then
- 10**  $R^6$  and  $R^7$  may also be selected from indanyl or indolyl, each of said indanyl or indolyl may be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkyloxycarbonyl, formyl, cyano, nitro, amino, and trifluoromethyl;
- 15**  $X^1$  and  $X^2$  are each independently  $-NR^3-$ ,  $-NH-NH-$ ,  $-N=N-$ ,  $-O-$ ,  $-S-$ ,  $-S(=O)-$  or  $-S(=O)_2-$ ;
- Alk is  $C_{1-4}$ alkanediyl; or
- when  $Y$  is  $CR^5$  then  $L$  may also be selected from  $C_{1-10}$ alkyl,  $C_{3-10}$ alkenyl,  $C_{3-10}$ alkynyl,  $C_{3-7}$ cycloalkyl, or  $C_{1-10}$ alkyl
- 20** substituted with one or two substituents independently selected from  $C_{3-7}$ cycloalkyl, indanyl, indolyl and phenyl, wherein said phenyl, indanyl and indolyl may be substituted with one, two, three, four or where possible five substituents each independently selected from halo, hydroxy,
- 25**  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, cyano, aminocarbonyl,  $C_{1-6}$ alkyloxycarbonyl, formyl, nitro, amino, trihalomethyl, trihalomethyloxy and  $C_{1-6}$ alkylcarbonyl;
- aryl is phenyl or phenyl substituted with one, two, three, four
- 30** or five substituents each independently selected from halo,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, cyano, nitro and trifluoromethyl;
- Het is an aliphatic or aromatic heterocyclic radical; said aliphatic heterocyclic radical is selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl,
- 35** tetrahydrofuranyl and tetrahydrothienyl wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic heterocyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein each of said
- 40** aromatic heterocyclic radical may optionally be substituted with hydroxy.

The compounds of formula (I) can be prepared according to the methods described in the patent applications with application

**45** number PCT/EP99/02043 and PCT/EP99/02044.

## 3

(II) is an antiviral compound of formula



10 the *N*-oxides, the pharmaceutically acceptable addition salts, quaternary amines and the stereochemically isomeric forms thereof, wherein

-b¹=b²-C(R²ᵃ)=b³-b⁴= represents a bivalent radical of formula

- 15
- CH=CH-C(R²ᵃ)=CH-CH= (b-1);
  - N=CH-C(R²ᵃ)=CH-CH= (b-2);
  - CH=N-C(R²ᵃ)=CH-CH= (b-3);
  - N=CH-C(R²ᵃ)=N-CH= (b-4);
  - N=CH-C(R²ᵃ)=CH-N= (b-5);
  - CH=N-C(R²ᵃ)=N-CH= (b-6);
  - 20 -N=N-C(R²ᵃ)=CH-CH= (b-7);

*q* is 0, 1, 2; or where possible *q* is 3 or 4;


*R*¹ is hydrogen, aryl, formyl, C₁-₆alkylcarbonyl, C₁-₆alkyl, C₁-₆alkyloxycarbonyl, C₁-₆alkyl substituted with formyl, C₁-₆alkylcarbonyl, C₁-₆alkyloxycarbonyl;

25 *R*²ᵃ is cyano, aminocarbonyl, mono- or di(methyl)aminocarbonyl, C₁-₆alkyl substituted with cyano, aminocarbonyl or mono- or di(methyl)aminocarbonyl, C₂-₆alkenyl substituted with cyano, or C₂-₆alkynyl substituted with cyano;

each *R*² independently is hydroxy, halo, C₁-₆alkyl optionally sub-

30 stituted with cyano or -C(=O)*R*⁶, C₃-₇cycloalkyl, C₂-₆alkenyl optionally substituted with one or more halogen atoms or cyano, C₂-₆alkynyl optionally substituted with one or more halogen atoms or cyano, C₁-₆alkyloxy, C₁-₆alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C₁-₆alkyl)amino,

35 polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>*p*</sub>*R*⁶, -NH-S(=O)<sub>*p*</sub>*R*⁶, -C(=O)*R*⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)*R*⁶, -C(=NH)*R*⁶ or a radical of formula



wherein each *A* independently is N, CH or CR⁶;

*B* is NH, O, S or NR⁶;

45 *p* is 1 or 2; and

*R*⁶ is methyl, amino, mono- or dimethylamino or polyhalomethyl;


- L is C<sub>1-10</sub>alkyl, C<sub>2-10</sub>alkenyl, C<sub>2-10</sub>alkynyl, C<sub>3-7</sub>cycloalkyl, whereby each of said aliphatic group may be substituted with one or two substituents independently selected from
- \* C<sub>3-7</sub>cycloalkyl,
  - 5 \* indolyl or isoindolyl, each optionally substituted with one, two, three or four substituents each independently selected from halo, C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, polyhalomethyl, polyhalomethyloxy and C<sub>1-6</sub>alkylcarbonyl,
  - 10 \* phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R<sup>2</sup>; or
- 15 L is -X-R<sup>3</sup> wherein  
 R<sup>3</sup> is phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R<sup>2</sup>; and
- 20 X is -NR<sup>1</sup>-, -NH-NH-, -N=N-, -O-, -C(=O)-, -CHOH-, -S-, -S(=O)- or -S(=O)<sub>2</sub>-;
- Q represents hydrogen, C<sub>1-6</sub>alkyl, halo, polyhaloC<sub>1-6</sub>alkyl or -NR<sup>4</sup>R<sup>5</sup>; and
- 25 R<sup>4</sup> and R<sup>5</sup> are each independently selected from hydrogen, hydroxy, C<sub>1-12</sub>alkyl, C<sub>1-12</sub>alkyloxy, C<sub>1-12</sub>alkylcarbonyl, C<sub>1-12</sub>alkyloxy-carbonyl, aryl, amino, mono- or di(C<sub>1-12</sub>alkyl)amino, mono- or di(C<sub>1-12</sub>alkyl)aminocarbonyl wherein each of the aforementioned C<sub>1-12</sub>alkyl groups may optionally and each individually be
- 30 substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxy-carbonyl, cyano, amino, imino, mono- or di(C<sub>1-6</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup>, aryl and Het; or
- 35 R<sup>4</sup> and R<sup>5</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-12</sub>alkyl)aminoC<sub>1-4</sub>alkylidene;
- Y represents hydroxy, halo, C<sub>3-7</sub>cycloalkyl, C<sub>2-6</sub>alkenyl optionally substituted with one or more halogen atoms, C<sub>2-6</sub>alkynyl
- 40 optionally substituted with one or more halogen atoms, C<sub>1-6</sub>alkyl substituted with cyano or -C(=O)R<sup>6</sup>, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxy-carbonyl, carboxyl, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup> or aryl;
- 45

- aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C<sub>1-6</sub>alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, cyano, nitro, polyhaloC<sub>1-6</sub>alkyl and polyhaloC<sub>1-6</sub>alkyloxy;
- 5 Het is an aliphatic or aromatic heterocyclic radical; said aliphatic heterocyclic radical is selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl wherein each of said
- 10 aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said aromatic heterocyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein each of said aromatic heterocyclic radical may optionally be substituted with hydroxy.

15

The compounds of formula (II) can be prepared according to the methods described in the US patent applications with application number 60/143962 and 60/107792.



- 20 (III) is an antiviral compound of formula



25

(III)

- a N-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein
- 30 -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- represents a bivalent radical of formula



n is 0, 1, 2, 3 or 4; and in case -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- is (a-1), then n may also be 5;

R<sup>1</sup> is hydrogen, aryl, formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyl substituted with formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyloxycarbonyl; and


40

each R<sup>2</sup> independently is hydroxy, halo, C<sub>1-6</sub>alkyl optionally substituted with cyano or -C(=O)R<sup>4</sup>, C<sub>3-7</sub>cycloalkyl, C<sub>2-6</sub>alkenyl optionally substituted with one or more halogen atoms or cyano, C<sub>2-6</sub>alkynyl optionally substituted with one or more halogen atoms or cyano, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino,

45

polyhalomethyl, polyhalomethyloxy, polyhalomethylthio,  
 $-S(=O)_pR^4$ ,  $-NH-S(=O)_pR^4$ ,  $-C(=O)R^4$ ,  $-NHC(=O)H$ ,  $-C(=O)NHNH_2$ ,  
 $-NHC(=O)R^4$ ,  $-C(=NH)R^4$  or a radical of formula

5



(c)

wherein each A independently is N, CH or  $CR^4$ ;

B is NH, O, S or  $NR^4$ ;

10

p is 1 or 2; and

$R^4$  is methyl, amino, mono- or dimethylamino or polyhalo-  
 methyl;

L is  $C_{4-10}$ alkyl,  $C_{2-10}$ alkenyl,  $C_{2-10}$ alkynyl,  $C_{3-7}$ cycloalkyl,  
 whereby each of said aliphatic group may be substituted with  
 one or two substituents independently selected from

15

\*  $C_{3-7}$ cycloalkyl,

\* indolyl or isoindolyl, each optionally substituted with  
 one, two, three or four substituents each independently  
 selected from halo,  $C_{1-6}$ alkyl, hydroxy,  $C_{1-6}$ alkyloxy,  
 cyano, aminocarbonyl, nitro, amino, polyhalomethyl, poly-  
 halomethyloxy and  $C_{1-6}$ alkylcarbonyl,

20

\* phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl,  
 wherein each of said aromatic rings may optionally be  
 substituted with one, two, three, four or five substi-  
 tuents each independently selected from the substituents  
 defined in  $R^2$ ; or

25

L is  $-X-R^3$  wherein

$R^3$  is phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazi-  
 nyl, wherein each of said aromatic rings may optionally  
 be substituted with two, three, four or five substituents  
 each independently selected from the substituents defined  
 in  $R^2$ ; and

30

X is  $-NR^1-$ ,  $-NH-NH-$ ,  $-N=N-$ ,  $-O-$ ,  $-C(=O)-$ ,  $-CHOH-$ ,  $-S-$ ,  
 $-S(=O)-$  or  $-S(=O)_2-$ ;


35 aryl is phenyl or phenyl substituted with one, two, three, four  
 or five substituents each independently selected from halo,  
 $C_{1-6}$ alkyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy, cyano, nitro, poly-  
 halo $C_{1-6}$ alkyl and polyhalo $C_{1-6}$ alkyloxy.

40 The compounds of formula (III) can be prepared according to the  
 methods described in the US patent application with application  
 number 60/107799.

45

(IV) is an antiviral compound of formula

5



(IV)

10

the pharmaceutically acceptable acid addition salts and the stereochemically isomeric forms thereof, wherein

R<sup>1</sup> and R<sup>2</sup> are each independently selected from hydrogen; hydroxy;

15

amino; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; Ar<sup>1</sup>; mono- or di(C<sub>1-6</sub>alkyl)amino; mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl; dihydro-2(3H)-furanone; C<sub>1-6</sub>alkyl substituted with one or two substituents each independently selected from amino, imino, aminocarbonyl, aminocarbonylamino, hydroxy, hydroxyc<sub>1-6</sub>alkyloxy, carboxyl, mono- or di(C<sub>1-6</sub>alkyl)amino, C<sub>1-6</sub>alkyloxycarbonyl and thienyl; or R<sup>1</sup> and R<sup>2</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-4</sub>-alkylidene;

25

R<sup>3</sup> is hydrogen, Ar<sup>1</sup>, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyl substituted with C<sub>1-6</sub>alkyloxycarbonyl; and R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are each independently selected from hydrogen, hydroxy, halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethyloxy;

30

L is C<sub>1-10</sub>alkyl; C<sub>3-10</sub>alkenyl; C<sub>3-10</sub>alkynyl; C<sub>3-7</sub>cycloalkyl; or L is C<sub>1-10</sub>alkyl substituted with one or two substituents independently selected from C<sub>3-7</sub>cycloalkyl; indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C<sub>1-6</sub>alkylcarbonyl; phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C<sub>1-6</sub>alkylcarbonyl; and,

35

40 Ar<sup>1</sup> is phenyl, or phenyl substituted with one, two or three substituents each independently selected from halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, nitro or trifluoromethyl; with the proviso that compounds (a) to (o)

45




Co. No.	Alk	R <sup>1</sup> /R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>
5	a	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	CH <sub>3</sub>	H	H	H
	b	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	NO <sub>2</sub>	H
	c	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	C <sub>6</sub> H <sub>5</sub>	H	H	H	H
	d	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	NO <sub>2</sub>	H	CH <sub>3</sub>	H
	e	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	NH <sub>2</sub>	H
10	f	4-(2-methylpropyl)phenylmethyl	H/H	H	H	CF <sub>3</sub>	H	H
	g	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	Cl	H
	h	4-(2-methylpropyl)phenylmethyl	H/H	H	H	H	H	H
	i	3,4-dimethoxyphenylmethyl	H/H	H	H	H	H	H
	j	2,3-dimethoxyphenylmethyl	H/H	H	H	H	H	H
15	k	3,4-diethoxyphenylmethyl	H/H	H	H	H	H	H
	l	2-(3,5-(1,1-dimethylethyl)-4-hydroxy-phenyl)ethyl	H/H	H	H	H	H	H
	m	2-(3,5-(1,1-dimethylethyl)-4-hydroxy-phenyl)ethyl	H/H	H	H	t-Bu	OH	t-Bu
	n	Phenylmethyl	H/H	H	CH <sub>3</sub>	H	H	H
	o	Phenylmethyl	H/H	H	H	H	H	H

20 are not included.

The compounds of formula (IV) can be prepared according to the methods described in EP-A-0834507.

25 (V) is an antifungal compound of formula



the *N*-oxide forms, the pharmaceutically acceptable acid addition salts and stereochemically isomeric forms thereof, wherein

*n* is zero, 1, 2 or 3;

40 *X* is N or CH;


each *R*<sup>1</sup> independently is halo, nitro, cyano, amino, hydroxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyloxy or trifluoromethyl;

*R*<sup>2</sup> is hydrogen; C<sub>3-7</sub>alkenyl; C<sub>3-7</sub>alkynyl, aryl; C<sub>3-7</sub>cycloalkyl; C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-4</sub>alkyloxy, C<sub>3-7</sub>cycloalkyl or aryl;

45

*R*<sup>3</sup> and *R*<sup>4</sup> each independently are hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-7</sub>cycloalkyl or aryl; or

R<sup>3</sup> and R<sup>4</sup> taken together form a bivalent radical -R<sup>3</sup>-R<sup>4</sup>- of formula:



10


wherein R<sup>5a</sup>, R<sup>5b</sup>, R<sup>5c</sup>, R<sup>5d</sup> each independently are hydrogen, C<sub>1-6</sub>alkyl or aryl; and

aryl is phenyl or phenyl substituted with one, two or three substituents selected from halo, nitro, cyano, amino, hydroxy,

15 C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyloxy or trifluoromethyl.


The compounds of formula (V) can be prepared according to the methods described in WO 99/02523.

20 (VI) is an apolipoprotein-B synthesis inhibitor of formula



30

the N-oxides, the stereochemically isomeric forms thereof, and the pharmaceutically acceptable acid addition salts, wherein A and B taken together form a bivalent radical of formula :



40 in the bivalent radicals of formula (a) and (b) the hydrogen atom may be replaced by C<sub>1-6</sub>alkyl; in the bivalent radicals of formula (c), (d), (e), (f), one or two hydrogen atoms may be replaced by C<sub>1-6</sub>alkyl;

R<sup>1</sup> is hydrogen, C<sub>1-6</sub>alkyl or halo;

45 R<sup>2</sup> is hydrogen or halo;

R<sup>3</sup> is hydrogen; C<sub>1-8</sub>alkyl; C<sub>3-6</sub>cycloalkyl; or C<sub>1-8</sub>alkyl substituted with hydroxy, oxo, C<sub>3-6</sub>cycloalkyl or aryl;

- Het is a heterocycle selected from the group consisting of pyridine; pyridine substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino or aryl; pyrimidine;
- 5 pyrimidine substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)-amino or aryl; tetrazole; tetrazole substituted with C<sub>1-6</sub>alkyl or aryl; triazole; triazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl,
- 10 hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)-amino; thiadiazole; thiadiazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)-amino; oxadiazole substituted with one or two substituents
- 15 selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino; imidazole; imidazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino; thiazole; thiazole substituted with one or
- 20 two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino; oxazole; oxazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino;
- 25 aryl is phenyl or phenyl substituted with C<sub>1-6</sub>alkyl or halo.

The heterocyclic radical "Het" is bound to the sulfur atom via a carbon atom.

- 30 The compounds of formula (VI) can be prepared according to the methods described in WO 96/13499.

The particles comprise the compounds of formula (I) to (VI) as a solid dispersion in a polymeric matrix, wherein the poly-

35 meric matrix is consisting of a homo- or copolymer of N-vinylpyrrolidone. Furthermore, the invention concerns a process for manufacturing of such particles and pharmaceutical dosage forms comprising such particles.

- 40 The compounds of formula (I) to (VI) contained in the particles show poor bio-availability.

In order to improve the dissolution characteristics the compounds are dispersed in a polymeric matrix, preferably by using a melt-

45 extrusion process.

## 11

EP-A 0 240 904 discloses a method for producing solid pharmaceutical forms by extrusion of polymer melts which contain active substances, using as polymers homo- or copolymers of N-vinylpyrrolidone.

5

EP-B 0 580 860 discloses a method for producing solid dispersions of drug substances in a polymeric matrix using a twin screw extruder.

- 10 It is an object of the present invention to provide rate-controlled pharmaceutical forms containing the aforementioned compounds.

We have found that this object is achieved by the particles

- 15 defined at the outset.

Preferred compounds according to the invention are:

4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;

- 20 4-[[2-[(cyanophenyl)amino]-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;

4-[[4-amino-5-chloro-6-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]-amino]benzonitrile;

4-[[5-chloro-4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]-amino]benzonitrile;

- 25 4-[[5-bromo-4-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]-amino]benzonitrile;

4-[[4-amino-5-chloro-6-[(4-cyano-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;

- 30 4-[[5-bromo-6-[(4-cyano-2,6-dimethylphenyl)amino]-2-pyrimidinyl]-amino]benzonitrile;

4-[[4-amino-5-chloro-6-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;

4-[[4-amino-5-bromo-6-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;

- 35 4-[[4-[(2,4,6-trimethylphenyl)amino]-1,3,5-triazin-2-yl]amino]benzonitrile;

4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-1,3,5-triazin-2-yl]-amino]benzonitrile;

- 40 4-[[4-[(2,6-dichlorophenyl)methyl]-6-(hydroxyamino)-1,3,5-triazin-2-yl]amino]benzonitrile;

1-[4-[4-[4-[[4-(2,4-difluorophenyl)-4-(1H-1,2,4-triazol-1-yl)methyl]-1,3-dioxolan-2-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-3-(1-methylethyl)-2-imidazolidinone;

45

## 12

(-)-[2S-[2alpha,4alpha(S\*)]]-4-[4-[4-[4-[[2-(4-chlorophenyl)-2-[[[(4-methyl-4H-1,2,4-triazol-3-yl)thio]methyl]-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-2,4-dihydro-2-(1-methyl-propyl)-3H-1,2,4-triazol-3-one,

- 5 a N-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof.

According to the present invention the term "rate-controlled" means that depending on the composition of the matrix the

- 10 particles can show instant release of the active ingredient or modified release (sustained release).

The compounds according to the invention are homogeneously dispersed in a polymer matrix consisting of a homopolymer of

15 N-vinylpyrrolidone or, preferably, a copolymer of N-vinylpyrrolidone. A preferred copolymer is a copolymer of N-vinylpyrrolidone and vinyl acetate, especially a copolymer obtained from 60% b.w. of NVP and 40% b.w. of vinylacetate.

- 20 The polymers show Fikentscher K values of from 17 to 90, preferably a K value of 30 (for the definition of the K value see "H. Fikentscher, Cellulose-Chemie" (1932), 58-64 and 71-74).

The polymeric matrix component is used in amounts of from 40 to

25 70, preferably of from 50 to 65% b.w. of the total weight of the particles.

- In a preferred embodiment of the invention the polymeric matrix further comprises a surfactant, preferably a surfactant with
- 30 a HLB-value of 10-18 (HLB: Hydrophilic Lipophilic Balance). Especially preferred surfactants are selected from the group consisting of low molecular weight polyoxyethylene polyoxypropylene block copolymers with a mean molecular weight of 1000 to 6000 g/mol, and hydrogenated castor oil which can be
- 35 modified with polyethylene glycol.

The amounts of surfactants used lies in the range of up to 20% b.w., preferably 5 to 15% b.w., of the particles.

- 40 In another preferred embodiment the matrix further comprises an organic carboxylic acid in amounts of up to 5% b.w. of the particles.

In another preferred embodiment of the invention the polymeric matrix further comprises hydroxypropyl methyl cellulose in

- 45 amounts of up to 25% b.w., preferably from 5 to 10% b.w..

## 13

The particles of the present invention are prepared as solid dispersions of the active compounds in a polymeric matrix. The term "solid dispersion" is well known in the art and means a dispersion consisting of solid components. Preferably solid  
5 dispersions are in the form of solid solutions wherein the active ingredients are molecularly dispersed in the polymeric matrix.

Such solid dispersion is preferably obtained by forming a homogeneous mixture of the components in the form of a melt,  
10 extruding said melt and shaping of the extrudate. The melting is effected by the input of thermal and/or mechanic energy.

Depending on the composition of the matrix, the melting takes place in the range of from 40°C to 190°C, preferably 50 to 150°C.  
15 The suitable temperature range depends on the glass transition temperature of the polymer component, the properties of the active ingredients and further additives. The optimal temperature range can be established by a few simple tests.

20 The mixing of the active substances with the polymer and additional components of the matrix can take place before or after the melting of the polymer. Preferably the process is solvent-free which means that no additional organic solvents or water are added.

25 The plastification and melting preferably can take place in an extruder, a kneader or a mixing reactor, preferably in an extruder having one or more screws which may rotate in the same direction or opposite directions, especially in a twin screw  
30 extruder. The latter can be operated with or without kneading elements, but use of kneading elements is preferred because mixing is better.

The still plastic material is extruded through a die or a breaker  
35 plate and then shaped into particles. This may be effected by milling and subsequent sieving the cooled extrudate. The preferred particle size for instant release forms lies in the range of from 0.5 to 1.5 mm.

40 The particles, optionally together with conventional pharmaceutically acceptable excipients, may be further processed to conventional pharmaceutical dosage forms such as tablets, pastilles, suppositories, or be packed in capsules.

45 It is possible and particularly advantageous to produce pharmaceutical forms with rate-controlled release and improved dissolution rates of the active ingredients. This was not to be

expected in view of the low solubility of the active ingredients in aqueous media.

#### Examples

5

#### General method

Powder mixes of the components were melt kneaded at 145°C for 5 min.. After cooling the solidified melts were ground and  
 10 sieved. The sieve fraction 0.5-1.5 mm was used for the dissolution tests.

The composition of the individual powder mixes is listed in Table 1.

15

Table 1

	Example No.	1	2	3	4	5	6
	Active ingredient <sup>1)</sup>	30	30	30	30	30	40
20	VP-VAC-copolymer <sup>2)</sup>	65	55	55	60	55	47,1
	Surfactant <sup>3)</sup>	5	15		5	5	4,3
	Citric acid				5		
	HPMC					10	8,6
	Surfactant <sup>4)</sup>			15			

25 <sup>1)</sup> 4-[[4-[2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]-benzonitrile

<sup>2)</sup> Kollidon® VA64, VP/VAC = 60/40, BASF Aktiengesellschaft

<sup>3)</sup> PEG-n-hydrogenated Castoroil

<sup>4)</sup> polyoxyethylene polyoxypropylene blockcopolymer, mean mol.  
 30 weight 4000 g/mol

The dissolution tests were carried out according to USP XXIII, paddle model, pH no change test, 0.1 M HCl, at 37°C, 100 rpm

35

40

45

## 15

The results are listed in Table 2.

Table 2: Dissolution Rates of particles according to examples 1-6

5

10

15

time [min]	Dissolution [%]				time [min]	Dissolution [%]	
	Ex. 1 (IR)	Ex. 2 (IR)	Ex. 3 (IR)	Ex. 4 (IR)		Ex. 5 (SR)	Ex. 6 (SR)
5	53	65	58	57	1		
10	73	86	88	82	2		
15	77	91	95	89	3		
20	81	91	96	93	4		
30	87	94	99	94	6		
60	92	93	96	94	8	96	95
120	93	94	97	95			
IR: Instant Release					SR: Sustained Release		

20 DSC-Measurements were performed with the formulations according to examples 1 to 6 in open pans (air) at temperatures of from 20 → 250°C, with a heating rate of 10°C per minute. There is no indication of crystalline drug substance in the solid dispersions.

25

30

35

40


45



## Claims

1. Rate-controlled release particles, comprising a compound of  
 5 formula I

10



a N-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein

15

Y is CR<sup>5</sup> or N;

A is CH, CR<sup>4</sup> or N;

n is 0, 1, 2, 3 or 4;

Q is -NR<sup>1</sup>R<sup>2</sup> or when Y is CR<sup>5</sup> then Q may also be hydrogen;

20

R<sup>1</sup> and R<sup>2</sup> are each independently selected from hydrogen, hydroxy, C<sub>1-12</sub>alkyl, C<sub>1-12</sub>alkyloxy, C<sub>1-12</sub>alkylcarbonyl, C<sub>1-12</sub>alkyloxycarbonyl, aryl, amino, mono- or di(C<sub>1-12</sub>alkyl)-amino, mono- or di(C<sub>1-12</sub>alkyl)aminocarbonyl wherein each of the aforementioned C<sub>1-12</sub>alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxy-C<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, aminocarbonyl, aminocarbonylamino, mono- or di(C<sub>1-6</sub>alkyl)amino, aryl and Het; or

25

R<sup>1</sup> and R<sup>2</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-12</sub>alkyl)aminoC<sub>1-4</sub>-alkylidene;

R<sup>3</sup> is hydrogen, aryl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyl substituted with C<sub>1-6</sub>alkyloxycarbonyl; and

35

each R<sup>4</sup> independently is hydroxy, halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, or when Y is CR<sup>5</sup> then R<sup>4</sup> may also represent C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl;

40

R<sup>5</sup> is hydrogen or C<sub>1-4</sub>alkyl;

L is -X<sup>1</sup>-R<sup>6</sup> or -X<sup>2</sup>-Alk-R<sup>7</sup> wherein


45

R<sup>6</sup> and R<sup>7</sup> each independently are phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy,

## 17

5 C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl-  
oxycarbonyl, formyl, cyano, nitro, amino, and tri-  
fluoromethyl; or when Y is CR<sup>5</sup> then R<sup>6</sup> and R<sup>7</sup> may  
also be selected from phenyl substituted with one,  
two, three, four or five substituents each indepen-  
dently selected from aminocarbonyl, trihalomethyloxy  
and trihalomethyl; or when Y is N then R<sup>6</sup> and R<sup>7</sup> may  
also be selected from indanyl or indolyl, each of  
said indanyl or indolyl may be substituted with one,  
two, three, four or five substituents each  
independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl,  
C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyloxycarbonyl,  
formyl, cyano, nitro, amino, and trifluoromethyl;  
10 X<sup>1</sup> and X<sup>2</sup> are each independently -NR<sup>3</sup>-, -NH-NH-, -N=N-,  
-O-, -S-, -S(=O)- or -S(=O)<sub>2</sub>-;  
15 Alk is C<sub>1-4</sub>alkanediyl; or  
when Y is CR<sup>5</sup> then L may also be selected from C<sub>1-10</sub>alkyl,  
C<sub>3-10</sub>alkenyl, C<sub>3-10</sub>alkynyl, C<sub>3-7</sub>cycloalkyl, or C<sub>1-10</sub>alkyl  
substituted with one or two substituents independently  
selected from C<sub>3-7</sub>cycloalkyl, indanyl, indolyl and  
20 phenyl, wherein said phenyl, indanyl and indolyl may be  
substituted with one, two, three, four or where possible  
five substituents each independently selected from halo,  
hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl,  
C<sub>1-6</sub>alkyloxycarbonyl, formyl, nitro, amino, trihalomethyl,  
25 trihalomethyloxy and C<sub>1-6</sub>alkylcarbonyl;  
aryl is phenyl or phenyl substituted with one, two, three,  
four or five substituents each independently selected  
from halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, nitro and  
trifluoromethyl;  
30 Het is an aliphatic or aromatic heterocyclic radical;  
said aliphatic heterocyclic radical is selected from  
pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl,  
morpholinyl, tetrahydrofuranyl and tetrahydrothienyl  
wherein each of said aliphatic heterocyclic radical may  
optionally be substituted with an oxo group; and said  
aromatic heterocyclic radical is selected from pyrrolyl,  
furanyl, thienyl, pyridyl, pyrimidinyl, pyrazinyl and  
pyridazinyl wherein each of said aromatic heterocyclic  
40 radical may optionally be substituted with hydroxy,

or a compound of formula



10 the *N*-oxides, the pharmaceutically acceptable addition salts, quaternary amines and the stereochemically isomeric forms thereof, wherein

$-b^1=b^2-C(R^{2a})=b^3-b^4=$  represents a bivalent radical of formula

$$-\text{CH}=\text{CH}-\text{C}(\text{R}^{2a})=\text{CH}-\text{CH}= \quad (\text{b-1}) ;$$

**15**  $-N=CH-C(R^{2a})=CH-CH=$  (b-2) ;

$$-\text{CH}=\text{N}-\text{C}(\text{R}^{2a})=\text{CH}-\text{CH}= \quad (\text{b-3}) ;$$
$$-N=CH-C(R^{2a})=N-CH= \quad (b-4) ;$$
$$-\text{N}=\text{CH}-\text{C}(\text{R}^{2\text{a}})=\text{CH}-\text{N}= \quad (\text{b-5}) ;$$
$$-\text{CH}=\text{N}-\text{C}(\text{R}^{2a})=\text{N}-\text{CH}= \quad (\text{b-6}) ;$$


20  $-N=N-C(R^{2a})=CH-CH=$  (b-7) ;

$q$  is 0, 1, 2; or where possible  $q$  is 3 or 4;

R<sup>1</sup> is hydrogen, aryl, formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyl substituted with formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyloxycarbonyl;

25 R<sup>2a</sup> is cyano, aminocarbonyl, mono- or di(methyl)amino-  
carbonyl, C<sub>1-6</sub>alkyl substituted with cyano, amino-  
carbonyl or mono- or di(methyl)aminocarbonyl, C<sub>2-6</sub>alkenyl  
substituted with cyano, or C<sub>2-6</sub>alkynyl substituted with  
cyano;

30 each R<sup>2</sup> independently is hydroxy, halo, C<sub>1-6</sub>alkyl optionally substituted with cyano or -C(=O)R<sup>6</sup>, C<sub>3-7</sub>cycloalkyl, C<sub>2-6</sub>alkenyl optionally substituted with one or more halogen atoms or cyano, C<sub>2-6</sub>alkynyl optionally substituted with one or more halogen atoms or cyano, C<sub>1-6</sub>alkyloxy, 35 C<sub>1-6</sub>alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup> or a radical of formula



45

## 19

wherein each A independently is N, CH or CR<sup>6</sup>;

B is NH, O, S or NR<sup>6</sup>;

p is 1 or 2; and

R<sup>6</sup> is methyl, amino, mono- or dimethylamino or polyhalomethyl;

5

L is C<sub>1-10</sub>alkyl, C<sub>2-10</sub>alkenyl, C<sub>2-10</sub>alkynyl, C<sub>3-7</sub>cycloalkyl, whereby each of said aliphatic group may be substituted with one or two substituents independently selected from

10

\* C<sub>3-7</sub>cycloalkyl,

\* indolyl or isoindolyl, each optionally substituted with one, two, three or four substituents each independently selected from halo, C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, polyhalomethyl, polyhalomethyloxy and C<sub>1-6</sub>alkyl-carbonyl,

15

\* phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R<sup>2</sup>; or

20

L is -X-R<sup>3</sup> wherein

R<sup>3</sup> is phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in R<sup>2</sup>; and

25

X is -NR<sup>1</sup>-, -NH-NH-, -N=N-, -O-, -C(=O)-, -CHOH-, -S-, -S(=O)- or -S(=O)<sub>2</sub>-;

Q represents hydrogen, C<sub>1-6</sub>alkyl, halo, polyhaloC<sub>1-6</sub>alkyl or -NR<sup>4</sup>R<sup>5</sup>; and

30

R<sup>4</sup> and R<sup>5</sup> are each independently selected from hydrogen, hydroxy, C<sub>1-12</sub>alkyl, C<sub>1-12</sub>alkyloxy, C<sub>1-12</sub>alkylcarbonyl, C<sub>1-12</sub>alkyloxycarbonyl, aryl, amino, mono- or di(C<sub>1-12</sub>alkyl)amino, mono- or di(C<sub>1-12</sub>alkyl)aminocarbonyl wherein each of the aforementioned C<sub>1-12</sub>alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, mono- or di(C<sub>1-6</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup>, aryl and Het; or

35


40

R<sup>4</sup> and R<sup>5</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-12</sub>alkyl)aminoC<sub>1-4</sub>-alkylidene;

45

## 20


- Y represents hydroxy, halo, C<sub>3-7</sub>cycloalkyl, C<sub>2-6</sub>alkenyl optionally substituted with one or more halogen atoms, C<sub>2-6</sub>alkynyl optionally substituted with one or more halogen atoms, C<sub>1-6</sub>alkyl substituted with cyano or
- 5 -C(=O)R<sup>6</sup>, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup> or aryl;
- 10 aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, C<sub>1-6</sub>alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, cyano, nitro, polyhaloC<sub>1-6</sub>alkyl and polyhaloC<sub>1-6</sub>alkyloxy;
- Het is an aliphatic or aromatic heterocyclic radical;
- 15 said aliphatic heterocyclic radical is selected from pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl and tetrahydrothienyl wherein each of said aliphatic heterocyclic radical may optionally be substituted with an oxo group; and said
- 20 aromatic heterocyclic radical is selected from pyrrolyl, furanyl, thienyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl wherein each of said aromatic heterocyclic radical may optionally be substituted with hydroxy,
- 25 or a compound of formula



- a N-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof,
- 35 wherein
- a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- represents a bivalent radical of formula
- CH=CH-CH=CH- (a-1);
- N=CH-CH=CH- (a-2);
- N=CH-N=CH- (a-3);
- 40 -N=CH-CH=N- (a-4);
- N=N-CH=CH- (a-5);
- n is 0, 1, 2, 3 or 4; and in case -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- is (a-1), then n may also be 5;
- R<sup>1</sup> is hydrogen, aryl, formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl,
- 45 C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyl substituted with formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyloxycarbonyl; and

## 21

each  $R^2$  independently is hydroxy, halo,  $C_{1-6}$ alkyl optionally substituted with cyano or  $-C(=O)R^4$ ,  $C_{3-7}$ cycloalkyl,  $C_{2-6}$ alkenyl optionally substituted with one or more halogen atoms or cyano,  $C_{2-6}$ alkynyl optionally substituted with one or more halogen atoms or cyano,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di( $C_{1-6}$ alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio,  $-S(=O)_pR^4$ ,  $-NH-S(=O)_pR^4$ ,  $-C(=O)R^4$ ,  $-NHC(=O)H$ ,  $-C(=O)NHNH_2$ ,  $-NHC(=O)R^4$ ,  $-C(=NH)R^4$  or a radical of formula




(c)

- wherein each A independently is N, CH or  $CR^4$ ;  
 B is NH, O, S or  $NR^4$ ;  
 p is 1 or 2; and  
 $R^4$  is methyl, amino, mono- or dimethylamino or polyhalomethyl;
- L is  $C_{4-10}$ alkyl,  $C_{2-10}$ alkenyl,  $C_{2-10}$ alkynyl,  $C_{3-7}$ cycloalkyl, whereby each of said aliphatic group may be substituted with one or two substituents independently selected from
- \*  $C_{3-7}$ cycloalkyl,
  - \* indolyl or isoindolyl, each optionally substituted with one, two, three or four substituents each independently selected from halo,  $C_{1-6}$ alkyl, hydroxy,  $C_{1-6}$ alkyloxy, cyano, aminocarbonyl, nitro, amino, polyhalomethyl, polyhalomethyloxy and  $C_{1-6}$ alkyl-carbonyl,
  - \* phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with one, two, three, four or five substituents each independently selected from the substituents defined in  $R^2$ ; or
- L is  $-X-R^3$  wherein
- $R^3$  is phenyl, pyridinyl, pyrimidinyl, pyrazinyl or pyridazinyl, wherein each of said aromatic rings may optionally be substituted with two, three, four or five substituents each independently selected from the substituents defined in  $R^2$ ; and
- X is  $-NR^1-$ ,  $-NH-NH-$ ,  $-N=N-$ ,  $-O-$ ,  $-C(=O)-$ ,  $-CHOH-$ ,  $-S-$ ,  $-S(=O)-$  or  $-S(=O)_2-$ ;
- aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo,  $C_{1-6}$ alkyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy, cyano, nitro, polyhalo $C_{1-6}$ alkyl and polyhalo $C_{1-6}$ alkyloxy,

or a compound of formula

5



(IV)

10

the pharmaceutically acceptable acid addition salts and the stereochemically isomeric forms thereof, wherein

R<sup>1</sup> and R<sup>2</sup> are each independently selected from hydrogen;

15

hydroxy; amino; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; Ar<sup>1</sup>; mono- or di(C<sub>1-6</sub>alkyl)amino; mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl; dihydro-2(3H)-furanone; C<sub>1-6</sub>alkyl substituted with one or two substituents each independently selected from amino, imino, aminocarbonyl, aminocarbonylamino, hydroxy, hydroxyc<sub>1-6</sub>alkyloxy, carboxyl, mono- or di(C<sub>1-6</sub>alkyl)amino, C<sub>1-6</sub>alkyloxycarbonyl and thienyl; or

20

R<sup>1</sup> and R<sup>2</sup> taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-4</sub>-alkylidene;

25

R<sup>3</sup> is hydrogen, Ar<sup>1</sup>, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkyl substituted with C<sub>1-6</sub>alkyloxycarbonyl; and

30

R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are each independently selected from hydrogen, hydroxy, halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethoxy ;

L is C<sub>1-10</sub>alkyl; C<sub>3-10</sub>alkenyl; C<sub>3-10</sub>alkynyl; C<sub>3-7</sub>cycloalkyl; or

35

L is C<sub>1-10</sub>alkyl substituted with one or two substituents independently selected from C<sub>3-7</sub>cycloalkyl; indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy, C<sub>1-6</sub>alkylcarbonyl; phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy, C<sub>1-6</sub>alkylcarbonyl; and,

45


## 23

Ar<sup>1</sup> is phenyl, or phenyl substituted with one, two or three substituents each independently selected from halo, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, cyano, nitro or trifluoromethyl; with the proviso that compounds (a) to (o)

Co. No.	Alk	R <sup>1</sup> /R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>
a	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	CH <sub>3</sub>	H	H	H	H
b	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	NO <sub>2</sub>	H	H
c	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	C <sub>6</sub> H <sub>5</sub>	H	H	H	H	H
d	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	NO <sub>2</sub>	H	CH <sub>3</sub>	H	H
e	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	NH <sub>2</sub>	H	H
f	4-(2-methylpropyl)phenylmethyl	H/H	H	H	CF <sub>3</sub>	H	H	H
g	1-(4-(2-methylpropyl)phenyl)ethyl	H/H	H	H	H	Cl	H	H
h	4-(2-methylpropyl)phenylmethyl	H/H	H	H	H	H	H	H
i	3,4-dimethoxyphenylmethyl	H/H	H	H	H	H	H	H
j	2,3-dimethoxyphenylmethyl	H/H	H	H	H	H	H	H
k	3,4-diethoxyphenylmethyl	H/H	H	H	H	H	H	H
l	2-(3,5-(1,1-dimethylethyl)-4-hydroxy-phenyl)ethyl	H/H	H	H	H	H	H	H
m	2-(3,5-(1,1-dimethylethyl)-4-hydroxy-phenyl)ethyl	H/H	H	H	t-Bu	OH	t-Bu	H
n	Phenylmethyl	H/H	H	CH <sub>3</sub>	H	H	H	H
o	Phenylmethyl	H/H	H	H	H	H	H	H

are not included,

or a compound of formula



the *N*-oxide forms, the pharmaceutically acceptable acid addition salts and stereochemically isomeric forms thereof, wherein

*n* is zero, 1, 2 or 3;

*X* is N or CH;

each R<sup>1</sup> independently is halo, nitro, cyano, amino, hydroxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyloxy or trifluoromethyl;


R<sup>2</sup> is hydrogen; C<sub>3-7</sub>alkenyl; C<sub>3-7</sub>alkynyl, aryl; C<sub>3-7</sub>cycloalkyl; C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-4</sub>alkyloxy, C<sub>3-7</sub>cycloalkyl or aryl;



## 24

R<sup>3</sup> and R<sup>4</sup> each independently are hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-7</sub>cycloalkyl or aryl; or  
 R<sup>3</sup> and R<sup>4</sup> taken together form a bivalent radical -R<sup>3</sup>-R<sup>4</sup>- of formula:


5



15 wherein R<sup>5a</sup>, R<sup>5b</sup>, R<sup>5c</sup>, R<sup>5d</sup> each independently are hydrogen, C<sub>1-6</sub>alkyl or aryl; and aryl is phenyl or phenyl substituted with one, two or three substituents selected from halo, nitro, cyano, amino, hydroxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyloxy or trifluoromethyl,

or a compound of formula

20



30 the *N*-oxides, the stereochemically isomeric forms thereof, and the pharmaceutically acceptable acid addition salts, wherein A and B taken together form a bivalent radical of formula :

- 35
- N=CH- (a),
  - CH=N- (b),
  - CH<sub>2</sub>-CH<sub>2</sub>- (c),
  - CH=CH- (d),
  - C(=O)-CH<sub>2</sub>- (e),
  - CH<sub>2</sub>-C(=O)- (f),

40 in the bivalent radicals of formula (a) and (b) the hydrogen atom may be replaced by C<sub>1-6</sub>alkyl; in the bivalent radicals of formula (c), (d), (e), (f), one or two hydrogen atoms may be replaced by C<sub>1-6</sub>alkyl;

R<sup>1</sup> is hydrogen, C<sub>1-6</sub>alkyl or halo;

45 R<sup>2</sup> is hydrogen or halo;

R<sup>3</sup> is hydrogen; C<sub>1-8</sub>alkyl; C<sub>3-6</sub>cycloalkyl; or C<sub>1-8</sub>alkyl substituted with hydroxy, oxo, C<sub>3-6</sub>cycloalkyl or aryl;

Het is a heterocycle selected from the group consisting of pyridine; pyridine substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino or aryl; pyrimidine; pyrimidine substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)-amino or aryl; tetrazole; tetrazole substituted with C<sub>1-6</sub>alkyl or aryl; triazole; triazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)-amino; thiadiazole; thiadiazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)-amino; oxadiazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino; imidazole; imidazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino; thiazole; thiazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino; oxazole; oxazole substituted with one or two substituents selected from C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxy, trihalomethyl, amino, mono- or di(C<sub>1-6</sub>alkyl)amino; aryl is phenyl or phenyl substituted with C<sub>1-6</sub>alkyl or halo, and the heterocyclic radical "Het" is bound to the sulfur atom via a carbon atom,

as a solid dispersion in a polymeric matrix, wherein the polymeric matrix is consisting of a homo- or copolymer of N-vinylpyrrolidone.

2. Particles according to claim 1, wherein the copolymer of N-vinylpyrrolidone is a copolymer with vinyl acetate.

3. Particles according to claim 1 or 2, further comprising a surfactant.

4. Particles according to claim 3, wherein the surfactant is a PEG-n-hydrogenated castor oil.

5. Particles according to any of the claims 1 to 3, wherein the surfactant is a low molecular weight polyoxyethylene polyoxypropylene block copolymer.

6. Particles according to any of the claims 1 to 3, further comprising citric acid in amounts of up to 5 % b.w.
7. Particles according to any of the claims 1 to 6, wherein the  
5 homo- or copolymer of N-vinylpyrrolidone is used in amounts of from 40 to 70 % b.w. of the total weight of the dosage form.
8. Particles according to claim 7, wherein the homo- or copolymer of N-vinylpyrrolidone is used in amounts of from 50 to 65  
10 % b.w..
9. Particles according to any of the claims 1 to 8, wherein the controlled release is an instant release of the drug.
10. Particles according to any of the claims 1 to 8, wherein the  
15 controlled release is a sustained release.
11. Particles according to claim 10, further comprising hydroxypropyl methyl cellulose in amounts of from 5 to 10 % b.w..  
20
12. Particles according to any of the claims 1 to 11, obtained by forming a homogeneous mixture of the components in the form of a melt, extruding said mixture and shaping of the extrudate.
13. Particles according to any of the claims 1 to 11, comprising  
25 a compound selected from the group consisting of  
4-[[4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
30 4-[[2-[(cyanophenyl)amino]-4-pyrimidinyl]amino]-3,5-dimethylbenzonitrile;  
4-[[4-amino-5-chloro-6-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]-amino]benzonitrile;  
4-[[5-chloro-4-[(2,4,6-trimethylphenyl)amino]-2-pyrimidinyl]-  
35 amino]benzonitrile;  
4-[[5-bromo-4-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]-amino]benzonitrile;  
4-[[4-amino-5-chloro-6-[(4-cyano-2,6-dimethylphenyl)amino]-2-pyrimidinyl]amino]benzonitrile;  
40 4-[[5-bromo-6-[(4-cyano-2,6-dimethylphenyl)amino]-2-pyrimidinyl]-amino]benzonitrile;  
4-[[4-amino-5-chloro-6-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;  
45 4-[[4-amino-5-bromo-6-(4-cyano-2,6-dimethylphenoxy)-2-pyrimidinyl]amino]benzonitrile;

## 27

- 4-[[4-[(2,4,6-trimethylphenyl)amino]-1,3,5-triazin-2-yl]-amino]benzonitrile;  
4-[[4-amino-6-[(2,6-dichlorophenyl)methyl]-1,3,5-triazin-2-yl]amino]benzonitrile;  
**5** 4-[[4-[(2,6-dichlorophenyl)methyl]-6-(hydroxyamino)-1,3,5-triazin-2-yl]amino]benzonitrile;  
1-[4-[4-[4-[4-(2,4-difluorophenyl)-4-(1H-1,2,4-triazol-1-yl-methyl)-1,3-dioxolan-2-yl]methoxy]phenyl]-1-piperazinyl]-phenyl]-3-(1-methylethyl)-2-imidazolidinone;  
**10** (-)-[2S-[2alpha,4alpha(S\*)]]-4-[4-[4-[4-[[2-(4-chlorophenyl)-2-[[4-methyl-4H-1,2,4-triazol-3-yl]thio]methyl]-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-2,4-dihydro-2-(1-methyl-propyl)-3H-1,2,4-triazol-3-one,  
a N-oxide, a pharmaceutically acceptable addition salt or a  
**15** stereochemically isomeric form thereof.
14. Pharmaceutical dosage form, comprising particles according to any of the preceding claims.
- 20** 15. Pharmaceutical dosage forms according to claim 13, further comprising one or more pharmaceutically acceptable excipients.

25

30

35

40

45